

## **Bioaccumulation and effect of sediment-associated silver in different forms in two marine deposit feeders**

Dai, Lina; Banta, Gary Thomas; Syberg, Kristian; Selck, Henriette; Gilliland, Douglas; Forbes, Valery E.

*Publication date:*  
2012

*Document Version*  
Early version, also known as pre-print

*Citation for published version (APA):*  
Dai, L., Banta, G. T., Syberg, K., Selck, H., Gilliland, D., & Forbes, V. E. (2012). *Bioaccumulation and effect of sediment-associated silver in different forms in two marine deposit feeders*.

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact [rucforsk@kb.dk](mailto:rucforsk@kb.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Bioaccumulation and effect of sediment-associated silver in different forms in two marine deposit feeders

Lina Dai<sup>1</sup>, Gary T Banta<sup>1</sup>, Kristian Syberg<sup>1</sup>, Henriette Selck<sup>1</sup>, Douglas Gilliland<sup>2</sup> and Valery E Forbes<sup>1,3</sup>  
<sup>1</sup>Dept. of Environmental, Social and Spatial Change, Roskilde University, Denmark.  
<sup>2</sup>European Commission, Joint Research Centre, Institute for Health and Consumer Protection, Italy.  
<sup>3</sup>School of Biological Sciences, University of Nebraska Lincoln, Lincoln.  
E-mail contact: [ldai@ruc.dk](mailto:ldai@ruc.dk)




### Introduction

- Different behaviour and effects of metal-bearing nanoparticles (NPs) have been found compared to their corresponding metallic ions [1,2].
  - Toxicity of metal-bearing NPs isn't easily predicted when comparing to corresponding ionic form
  - It is unclear whether toxicity of metal-bearing NPs is dependent on particle size.
- The aim of our study is to exam effects at the individual level by measuring typical endpoints in two organisms (i.e., a marine polychaete, *Capitella teleta* and a marine bivalve, *Macoma balthica*) after exposure to sediment amended with different forms and particle sizes of Ag.

### Hypothesis:

Toxicity and biota is metal form/particle size dependent?  
(Here are their relative differences in size)



### Result – *Macoma balthica*

#### Toxicity

No negative effects were detected on mortality, condition index or growth of exposed clams for any Ag form (data not shown).

#### Bioaccumulation

Bioaccumulation of Ag in *M. balthica* decreased significantly with increasing particle size (One-way ANOVA,  $p = 0.03$ ) (Figure 1).





Figure 1. Silver concentration measured in *M. balthica* exposed to a nominal conc. of 200µg/g dw sed. \*\*\* refers to a significant difference from ionic Ag. Error bars indicated 1 standard deviation (n=5).

Reference  
[1] García-Alonso J, Khan FR, Misra SK, Turmaine M, Smith BD, Rainbow PS, Luoma SN, Valsami-Jones E. 2011. Cellular Internalization of Silver Nanoparticles in Gut Epithelia of the Estuarine Polychaete *Nereis diversicolor*. *Environ Sci Technol* 45:4630-4636. [2] Cong Y, Banta GT, Selck H, Berhanu D, Valsami-Jones E, Forbes VE. 2011. Toxic effects and bioaccumulation of nano-, micron- and ionic-Ag in the polychaete, *Nereis diversicolor*. *Aquatic Toxicology* 105:403-411. [3] Cong Y. 2011. PhD thesis of 'Toxic effects and bioaccumulation of nano-, micron- and aqueous-Ag in the estuarine polychaete, *Nereis (Hediste) diversicolor*', Roskilde University.

### Experimental design



Endpoints • Mortality • Growth • Health condition • Body burden

### Result – *Capitella teleta*

#### Toxicity

No significant effects on either mortality or specific growth rate were detected for any Ag form or nominal concentration (data not shown).

#### Bioaccumulation

There was no significant effect of Ag form on Ag accumulation in *C. teleta*, although body burden increased significantly as a function of nominal concentration (One-way ANOVA,  $p < 0.001$ ) (Figure 2).




Figure 2. Silver concentration measured in *Capitella teleta* exposed to nominal concentration of 10, 50 and 100 µg/g dw sed. Error bars indicated 1 standard deviation (n=4). 20 nm-nanoparticle at 100µg/g dw sed nominal concentration was removed due to a significant difference in initial measured Ag concentration from the other treatments with the same nominal concentration at day 0.

### Conclusions

- No significant effects on mortality and growth of *C. teleta* and *M. balthica*.
- All Ag forms are bioavailable to both organisms.
- Metal form/particle size dependence of bioavailability is species specific, possibly due to differences in:
  - gut structure, thus
  - particle sorting mechanisms

Such differences in the bioavailability of metal-bearing particles warrant further investigation and consideration in terms of the impact of them in sediment environments.